

Calanthe yuksomnensis (Orchidaceae: Epidendroideae): range extension and new addition to the flora of Meghalaya, Northeast India^a

Durgesh Verma^{1*} & Seshu Lavania²

Mots-clés/Keywords: *Calanthe*, endemism/endémicité, new addition/nouvel enregistrement, Meghalaya, NE India/Inde du nord-est.

Abstract

Calanthe yuksomnensis S.Z.Lucksom, an endemic species of India, is reported here as an addition to the orchid flora of the Meghalaya state. Its detailed description and photographic illustration are provided to facilitate identification.

Résumé

Calanthe yuksomnensis (Orchidaceae : Epidendroideae) : extension de sa distribution, nouvelle addition à la flore du Meghalaya, Inde du nord-est – *Calanthe yuksomnensis* S.Z.Lucksom, une espèce endémique d'Inde, est ici ajoutée à la flore d'orchidées de l'État du Meghalaya. Nous en proposons une description détaillée et une illustration afin de faciliter son identification.

Introduction

The genus *Calanthe* R.Brown (Orchidaceae) with ca. 209 species (Clayton & Cribb, 2013; Oh *et al.*, 2015) is distributed in tropical and subtropical Asia, Japan, SE Asia, New Guinea, Australia, Pacific Islands, tropical and South Africa, Madagascar, islands of the Indian Ocean, and the neotropics. In

^a : manuscrit reçu le 4 décembre 2015, accepté le 3 mars 2016

article mis en ligne sur www.richardiana.com le 05/03/2016 – pp. 189-194 - © Tropicalia
ISSN 1626-3596 (imp.) - 2262-9017 (élect.)

India it is represented by 33 species (Misra, 2007; Kumar *et al.*, 2014), with 9 species in Meghalaya (Chowdhery, 2009). *Calanthe* is characterized by plicate leaves, persistent floral bracts, and lip adnate to the column forming a tube with the spurred base (Clayton & Cribb, 2013).

During floristic exploration in Tuber sacred grove located in East Jaintia Hills District of Meghalaya, northeastern India, the senior author came across a curious plant belonging to the genus *Calanthe* which had not been previously reported from the state of Meghalaya. After a careful examination of morphological characters of the species and review of the pertinent literature (Hooker, 1890; King & Pantling, 1898; Kataki, 1986; Pearce & Cribb, 2002; Lucksom, 2007), it has been found that this material belongs to the endemic species *Calanthe yuksommensis* S.Z.Lucksom, hitherto known from Darjeeling and Sikkim, its type locality, but unrecorded from Meghalaya. This discovery widens the range of distribution of the species into Northeast India. The paper provides its detailed description and a photograph for future reference and correct identification.

Taxonomic description

Calanthe yuksommensis S.Z.Lucksom, *Journal of the Bombay Natural History Society* 95: 319, f. 1-8 (1998); Pearce N.R. & P.J.Cribb, *The orchids of Bhutan* 194-195 (2002); Lucksom S.Z., *The orchids of Sikkim and north east Himalaya* 428, f. 269 (2007); Yonzon R. *et al.*, *Life Science Bulletin* 9(2): 359-360 (2012); Clayton D. & P.Cribb, *The genus Calanthe* 152-153 (2013).

Type: India, Sikkim, Yuksom, 20.05.1996, Lucksom 311A (*holo.* CAL, *iso.* Gangtok, Forest Dept. Herb.).

Terrestrial herb, evergreen, 18-23 cm tall; pseudobulbs 1.3-1.8 × 1.0-1.5 cm, ovoid, with 2-4 annular rings; leaves 2-3, shortly petiolate, lamina 11.0-20.0 × 3.3-4.5 cm, elliptic-lanceolate, acute-acuminate, petiole channelled, 1.5-2.0 cm long; inflorescence solitary, synanthous, 5-7 flowered, peduncle 11-13 cm long, erect, puberulous, light-green with purplish tinge, base enclosed by 2-3 young leaves; peduncle & young leaves are enclosed by 2-3 unequal sheathing bracts, 3.0-5.5 cm long, many nerved, pale-green coloured when young and becomes fibrous and turns reddish-brown on maturity; peduncular bract 1, 0.7-0.8 cm long, ovate, acuminate, pale yellow, rachis terete, puberulous, 4-5 cm long; floral bracts 0.55-0.60 × ca. 0.25 cm, lanceolate, dull white with greenish tinge, 1-nerved; pedicellate

ovary 1.3-1.5 cm long, slightly curved, pubescent, pale-green with purplish tinge; flowers 3.1-3.7 cm across, whitish-brown, fragrant; dorsal sepal 1.4-1.5 × 0.70-0.75 cm, elliptic, acute, externally pubescent, brownish-orange with green tinge, 5-nerved; lateral sepals similar to dorsal sepal, spreading; petals 1.2-1.3 × 0.40-0.45 cm, rhombic, subacuminate, similar in color as sepals, 3-nerved, spreading; lip 1.2-1.3 × ca. 1.3 cm, 3-lobed, spurred, base clawed and adnate to column, lateral lobes oblong-ovate, obtuse, mid-lobe subreniform, 2-lobulate, lobules subobovate, obtuse, crenulated, disc with 3 shallow elongated calli; spur 1.9-2.1 cm long, slightly curved, cylindrical, pubescent; column ca. 7 mm long, hairy inside; anther ca. 3.5 mm in diameter, ovate; pollinia 8, obovate-clavate, subequal, yellow, ca. 3 mm long, attached to a translucent disc; fruit not seen. Fig. 1.

Flowering: March-April.

Habitat: the species was found growing in subtropical evergreen forest in association with *Begonia josephi* A.de Candolle, *Podocarpus nerifolius* D.Don, *Strobilanthes* and *Tectaria* sp.

Specimen examined: INDIA, Meghalaya, East Jaintia hills, Tuber sacred grove, 25°26'05.20"N & 92°16'23.28"E, 1359 m, 15.04.2014, *D.Verma* 314 (LWU).

Distribution: endemic to India [Sikkim: Yuksom and Tendong; Darjeeling: Dhorey and Rombi forest, Sukiapokhari; Meghalaya: Tuber, East Jaintia hills (present report)].

Note: the plants collected by the senior author from Meghalaya are characterised in having a shorter petiole than the Sikkimese plants from type locality.

Acknowledgements

The senior author is thankful to the Director, Botanical Survey of India (BSI), Kolkata and to the Head of Office, BSI, Northern Regional Centre, Dehradun, for the facilities and encouragement. Sincere thanks are also extended to Dr. Avishek Bhattacharjee, Scientist, Botanical Survey of India, Central National Herbarium, Howrah for his helpful suggestions, and to the Forest Department of Meghalaya for rendering logistic support during field exploration.




Fig. 1: *Calanthe yuksommensis* S.Z.Lucksom
(see opposite page)

Opposite page

Fig. 1: *Calanthe yuksommensis* S.Z.Lucksom

A. Habit (in situ), B. Vegetative shoot, C. Pseudobulbs, D. Leaf (adaxial surface), E. Floral bract, F. Flower (front view), G. Flower (back view), H. Dorsal sepal, I. Lateral sepals, J. Petals, K. Labellum, column, pedicel & ovary (dorsal view), L. Labellum, column, pedicel and ovary (lateral view), M. Pedicellate ovary, N. Labellum (adaxial view), O. Labellum (abaxial view), P. Column, Q. Anther cap (adaxial view), R. Anther cap (abaxial view), S. Pollinarium [A, not to scale]. Photographs by D.Verma

References

- Chowdhery, H.J., 2009. Orchid Diversity in North Eastern States of India. *Journal of the Orchid Society of India*. 23(1&2): 19-42.
- Clayton, D. & P.Cribb, 2013. *The genus Calanthe*. Natural History Publications (Borneo) Sdn. Bhd.: Kota Kinabalu and Royal Botanic Gardens, Kew. 399 pp.
- Hooker, J.D., 1890. Orchideae. In *Flora of British India* vol. 5: 667-858. L.Reeve & Co. London.
- Kataki, S.K., 1986. *Orchids of Meghalaya*. Forest Department, Shillong.
- King, G. & R.Pantling, 1898. The orchids of the Sikkim-Himalaya. *Annals of the Royal Botanic Garden, Calcutta* 8: 1-342.
- Kumar, R., A.Bhattacharjee, S.K.Singh & S.Sharma, 2014. A rare species of *Calanthe* R.Br. recorded for the first time from India. *Telopea* 16: 213-217.
- Lucksom, S.Z., 2007. *The orchids of Sikkim and north east Himalaya*. Spectrum House, Siliguri. 984 pp.
- Misra, S., 2007. *Orchids of India – A Glimpse*. Bishen Singh Mahendra Pal Singh, Dehradun. 402 pp.
- Oh, S.-H., H.-J.Suh & C.-W.Park, 2015. Two new species of *Calanthe* (Orchidaceae: Epidendroideae) from Korea. *Phytotaxa* 218(1): 69-76.
- Pearce, N.R. & P.J.Cribb, 2002. *The orchids of Bhutan*. Royal Botanic Garden Edinburgh & Royal Government of Bhutan. 643 pp.

1 : Botanical Survey of India, Northern Regional Centre, 192 Kaulagarh Road, Dehradun—248195, Uttarakhand, India.

* corresponding author - durgesh.kv@gmail.com

2 : Department of Botany, University of Lucknow, Lucknow—226007, Uttar Pradesh, India.

Nouvelles espèces de Pleurothallidinae (Orchidaceae) d'Espírito Santo, Brésil^a

Guy R. Chiron^{1*}, Renato Ximenes Bolsanello² & Fabio Tesch³

Mots-clés/Keywords/Palavras-chave : *Acianthera*, forêt atlantique brésilienne/
Brazilian atlantic forest/Mata Atlântica, *Pabstiella*, Serra do Castelo,
taxinomie/taxonomy/ taxonomia.

Résumé

Sont proposées dans cet article six nouvelles espèces de Pleurothallidinae originaires des forêts humides de l'État d'Espírito Santo, Brésil. L'une appartient au genre *Acianthera*, les autres au genre *Pabstiella*. Parmi ces dernières l'une possède des fleurs à sépales pubescents sur leur face interne tandis que les quatre autres produisent des fleurs entièrement glabres. Chaque espèce est décrite et illustrée et comparée aux taxons les plus apparentés.

Abstract

New species in Pleurothallidinae (Orchidaceae) from Espírito Santo, Brazil – Six new species of Pleurothallidinae collected in the rain forests of the State of Espírito Santo, Brazil, are here proposed. One of them belongs to the genus *Acianthera* whereas the other ones are members of *Pabstiella*. Among them one produces flowers with sepals pubescent on the internal side whereas the four other ones has flowers fully glabrous. Each species is described, illustrated and compared to their closest relatives.

Resumo

Novas espécies de Pleurothallidinae (Orchidaceae) do Espírito Santo, Brasil – Seis novas espécies de Pleurothallidinae coletadas na Mata Atlântica do Estado do Espírito Santo, Brasil, estão aqui propostas. Uma delas pertence ao gênero *Acianthera* enquanto as outras fazem parte do

^a : manuscrit reçu le 26 février 2016, accepté le 14 mars 2016

article mis en ligne sur www.richardiana.com le 15/03/2015 – pp. 194-214 - © Tropicalia
ISSN 1626-3596 (imp.) - 2262-9017 (élect.)